

**BAB VII**  
**DAFTAR PUSTAKA**

1. Noncommunicable diseases [Internet]. [cited 2019 Sep 7]. Available from: <https://www.who.int/news-room/fact-sheets/detail/noncommunicable-diseases>
2. Kemenkes RI. Hasil Utama Riskesdas 2018. 2018;
3. Dharmayanti M. IDAI - Overview adolescent health problems and services [Internet]. 2015 [cited 2019 Oct 31]. Available from: <http://www.idai.or.id/artikel/seputar-kesehatan-anak/overview-adolescent-health-problems-and-services>
4. Lang JJ, Phillips EW, Orpana HM, Tremblay MS, Ross R, Ortega FB, et al. Field-based measurement of cardiorespiratory fitness to evaluate physical activity interventions. Vol. 96, Bulletin of the World Health Organization. World Health Organization; 2018. p. 794–6.
5. Lim JU, Lee JH, Kim JS, Hwang Y Il, Kim TH, Lim SY, et al. Comparison of World Health Organization and Asia-Pacific body mass index classifications in COPD patients. *Int J COPD*. 2017 Aug 21;12:2465–75.
6. Cristina T, Lins B, Valente LM, Celestino D, Filho S, Barbosa O. testing and body mass index  $\mathcal{E}$ . *Rev Port Cardiol (English Ed [Internet]*. 2015;34(1):27–33. Available from: <http://dx.doi.org/10.1016/j.repce.2014.07.004>
7. Papathanasiou G, Georgakopoulos D, Papageorgiou E, Zerva E, Michalis L, Kalfakakou V, et al. Effects of smoking on heart rate at rest and during exercise, and on heart rate recovery, in young adults. *Hell J Cardiol*. 2013 May;54(3):168–77.
8. Maggioni AP, Maseri A, Fresco C, Franzosi MG, Mauri F, Santoro E, et al. The New England Journal of Medicine Downloaded from [nejm.org](http://nejm.org) at UC SHARED JOURNAL COLLECTION on February 14, 2011. For personal use only. No other uses without permission. Copyright © 1993 Massachusetts Medical Society. All rights reserved. *N Engl J Med*. 1993;329(20):1442–8.
9. Gonzaga LA, Vanderlei LCM, Gomes RL, Valenti VE. Caffeine affects autonomic control of heart rate and blood pressure recovery after aerobic exercise in young adults: A crossover study. *Sci Rep*. 2017 Dec 1;7(1).
10. Alberto Neder J, Nery LE, Peres C, Whipp BJ. Reference values for dynamic responses to incremental cycle ergometry in males and females aged 20 to 80. *Am J Respir Crit Care Med [Internet]*. 2001 Oct 15 [cited 2020 Nov 28];164(8 I):1481–6.

- Available from: <https://pubmed.ncbi.nlm.nih.gov/11704600/>
11. Hub XPTC. W. Larry Kenney, Jack Wilmore, David Costill-Physiology of Sport and Exercis [Internet]. [cited 2020 Nov 28]. Available from: [https://www.academia.edu/36989983/W\\_Larry\\_Kenney\\_Jack\\_Wilmore\\_David\\_Costill\\_Physiology\\_of\\_Sport\\_and\\_Exercis](https://www.academia.edu/36989983/W_Larry_Kenney_Jack_Wilmore_David_Costill_Physiology_of_Sport_and_Exercis)
  12. Sandbakk Ø, Ettema G, Leirdal S, Holmberg HC, Lacour JR. Gender differences in the physiological responses and kinematic behaviour of elite sprint cross-country skiers. Vol. 112, *European Journal of Applied Physiology*. 2012. p. 1087–94.
  13. Powell KE, Paluch AE, Blair SN. Physical Activity for Health: What Kind? How Much? How Intense? On Top of What? *Annu Rev Public Heal* [Internet]. 2011 [cited 2019 Sep 5];32:349–65. Available from: [www.annualreviews.org](http://www.annualreviews.org)
  14. HHS. Physical Activity Guidelines for Americans 2 nd edition.
  15. WHO | What is Moderate-intensity and Vigorous-intensity Physical Activity? WHO. 2014;
  16. WHO | Middle East respiratory syndrome coronavirus (MERS-CoV). WHO [Internet]. 2020 [cited 2020 Oct 29]; Available from: <http://www.who.int/emergencies/mers-cov/en/>
  17. Nicaise V, Crespo NC, Marshall S. Agreement between the IPAQ and accelerometer for detecting intervention-related changes in physical activity in a sample of latina women. *J Phys Act Heal*. 2014;11(4):846–52.
  18. Medina C, Barquera S, Janssen I. Validity and reliability of the International Physical Activity Questionnaire among adults in Mexico. *Rev Panam Salud Publica/Pan Am J Public Heal*. 2013;34(1):21–8.
  19. Burgomaster KA, Hughes SC, Heigenhauser GJF, Bradwell SN, Gibala MJ, Appl J, et al. Six sessions of sprint interval training increases muscle oxidative potential and cycle endurance capacity in humans Six sessions of sprint interval training increases muscle oxida-tive potential and cycle endurance capacity in humans. *J Appl Physiol* [Internet]. 1985 [cited 2019 Dec 15];98. Available from: <http://www.jap.org>
  20. Hambrecht R, Niebauer J, Marburger C, Grunze M, Kälberer B, Hauer K, et al. Various intensities of leisure time physical activity in patients with coronary artery disease: Effects on cardiorespiratory fitness and progression of coronary atherosclerotic lesions. *J Am Coll Cardiol*. 1993;22(2):468–77.
  21. Shepherd JT. Circulatory response to exercise in health. *Circulation* [Internet]. 1987 Dec [cited 2019 Nov 11];76(6 Pt 2):VI3-10. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/3315298>

22. Fagard RH. Exercise is good for your blood pressure: Effects of endurance training and resistance training. In: *Clinical and Experimental Pharmacology and Physiology*. 2006. p. 853–6.
23. Breisch EA, White FC, Nimmo LE, McKirnan MD, Bloor CM. Exercise-induced cardiac hypertrophy: A correlation of blood flow and microvasculature. *J Appl Physiol*. 1986;60(4):1259–67.
24. Moore RL, Musch TI, Yelamarty R V, Scaduto RC, Semanchick AM, Elensky M, et al. Chronic exercise alters contractility and morphology of isolated rat cardiac myocytes. *Am J Physiol [Internet]*. 1993 May [cited 2019 Nov 14];264(5 Pt 1):C1180-9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/8498479>
25. Buick FJ, Gledhill N, Froese AB, Spriet L, Meyers EC. Effect of induced erythrocythemia on aerobic work capacity. *J Appl Physiol Respir Environ Exerc Physiol*. 1980;48(4):636–42.
26. Bonsignore MR, Morici G, Santoro A, Pagano M, Cascio L, Bonanno A, et al. Circulating hematopoietic progenitor cells in runners. *J Appl Physiol*. 2002 Nov 1;93(5):1691–7.
27. Möbius-Winkler S, Uhlemann M, Adams V, Sandri M, Erbs S, Lenk K, et al. Coronary Collateral Growth Induced by Physical Exercise: Results of the Impact of Intensive Exercise Training on Coronary Collateral Circulation in Patients with Stable Coronary Artery Disease (EXCITE) Trial. *Circulation*. 2016 Apr 12;133(15):1438–48.
28. Oja P, Laukkanen R, Pasanen M, Tyry T, Vuori I. A 2-km walking test for assessing the cardiorespiratory fitness of healthy adults. *Int J Sports Med*. 1991;12(4):356–62.
29. Piccinno A, Colella D. Differences in cardiovascular fitness of Italian high-school adolescents according to different physical activity levels assessed by IPAQ-A: a cross-sectional study. *Sport Sci Health*. 2017;13(1):149–55.
30. Shook RP, Gribben NC, Hand GA, Paluch AE, Welk GJ, Jakicic JM, et al. Subjective estimation of physical activity using the international physical activity questionnaire varies by fitness level. *J Phys Act Heal*. 2016 Jan 1;13(1):79–86.
31. Steene-Johannessen J, Anderssen SA, Van Der Ploeg HP, Hendriksen IJM, Donnelly AE, Brage S, et al. Are self-report measures able to define individuals as physically active or inactive? *Med Sci Sports Exerc*. 2016;48(2):235–44.
32. Melanson EL, Freedson PS. Physical Activity Assessment: A Review of Methods

- [Internet]. Vol. 36, Critical Reviews in Food Science and Nutrition. Taylor and Francis Inc.; 1996 [cited 2020 Oct 17]. p. 385–96. Available from: <https://pubmed.ncbi.nlm.nih.gov/8725670/>
33. Shephard RJ. Limits to the measurement of habitual physical activity by questionnaires [Internet]. Vol. 37, British Journal of Sports Medicine. BMJ Publishing Group; 2003 [cited 2020 Oct 17]. p. 197–206. Available from: <https://pubmed.ncbi.nlm.nih.gov/12782543/>

