

DAFTAR PUSTAKA

1. Ahti T (2010) Risk Factors of Varicose Veins. Available at: <http://acta.uta.fi/pdf/978-951-44-8022-5.pdf>.
2. Baldt, M. M. et al. (1996) ‘Deep venous thrombosis of the lower extremity: Efficacy of spiral CT venography compared with conventional venography in diagnosis’, Radiology, 200(2), pp. 423–428. doi: 10.1148/radiology.200.2.8685336.
3. Barros Junior, N. de et al. (2010) ‘Gestação e varizes de membros inferiores: prevalência e fatores de risco’, Jornal Vascular Brasileiro, 9(2), pp. 29–35. doi: 10.1590/s1677-54492010000200004.
4. Beaglehole, R. et al. (1975) ‘Varicose veins in the South Pacific’, International Journal of Epidemiology, 4(4), pp. 295–299. doi: 10.1093/ije/4.4.295.
5. Beebe-Dimmer, J. L. et al. (2005) ‘The epidemiology of chronic venous insufficiency and varicose veins’, Annals of Epidemiology, 15(3), pp. 175–184. doi: 10.1016/j.annepidem.2004.05.015.
6. Bradbury, A. et al. (1999) ‘What are the symptoms of varicose veins? Edinburgh vein study cross sectional population survey’, British Medical Journal, 318(7180), pp. 353–356. doi: 10.1136/bmj.318.7180.353.
7. Brand, F. N. et al. (1988) ‘The epidemiology of varicose veins: The Framingham Study’, American Journal of Preventive Medicine, pp. 96–101. doi: 10.1016/s0749-3797(18)31203-0.

8. Callam, M. J. (1994) ‘Epidemiology of varicose veins’, *British Journal of Surgery*, 81(2), pp. 167–173. doi: 10.1002/bjs.1800810204.
9. Chen, C. L. and Guo, H. R. (2014) ‘Varicose veins in hairdressers and associated risk factors: A cross-sectional study’, *BMC Public Health*, 14(1). doi: 10.1186/1471-2458- 14-885.
10. Evans, C. J. et al. (1999) ‘Prevalence of varicose veins and chronic venous insufficiency in men and women in the general population: Edinburgh Vein Study’, *Journal of Epidemiology and Community Health*, 53(3), pp. 149–153. doi: 10.1136/jech.53.3.149.
11. Hamdan, A. (2012) ‘Management of varicose veins and venous insufficiency’, *JAMA - Journal of the American Medical Association*, 308(24), pp. 2612–2621. doi: 10.1001/jama.2012.111352.
12. Horrocks, E., Roake, J. and Lewis, D. (2008) ‘Best practice for assessment of patients with varicose veins’, *New Zealand Medical Journal*.
13. Jacobs, B. N. et al. (2017) ‘Pathophysiology of varicose veins’, *Journal of Vascular Surgery: Venous and Lymphatic Disorders*, 5(3), pp. 460–467. doi: 10.1016/j.jvsv.2016.12.014.
14. Kohno, K. et al. (2014) ‘Standing posture at work and overweight exacerbate varicose veins: Shimane CoHRE study’, *Journal of Dermatology*, 41(11), pp. 964–968. doi: 10.1111/1346-8138.12643.

15. Lee, A. J. et al. (2003) ‘Lifestyle factors and the risk of varicose veins: Edinburgh Vein Study’, *Journal of Clinical Epidemiology*, 56(2), pp. 171–179. doi: 10.1016/S0895-4356(02)00518-8.
16. Naoum, J. J. and Hunter, G. C. (2007) ‘Pathogenesis of varicose veins and implications for clinical management’, *Vascular*, 15(5), pp. 242–249. doi: 10.2310/6670.2007.00069.
17. WL Miller Chronic venous insufficiency *Cardiovasc Clin*, 22 (1992), pp. 67-100.
18. Enrique Criado M.D.(Clinical Instructor) George J.Jr.M.D.(Roscoe B.G. Cowper Professor of Surgery, Coordinator of General Surgery Services, Director of Vascular Surgery, Vice Chairman) ‘Venous disease’ doi : 10.1016/0011-3840(91)90002-7
19. Thayer E.S, Wayne W, La MorteMD, PhD, MPH Daniel R.G.MD James O.MMD Risk factors for chronic venous insufficiency: A dual case-control study. Doi : 10.1016/S0741-5214(95)70050-1