

BIBLIOGRAPHY

- [AOAC]. Association of Official Analytical Chemists. (2005). *Official Methods of Analysis of the Association of Official Analytical Chemistry*. Washington DC: AOAC, Inc.
- [ITIS]. Integrated Taxonomic Information System. (2011). *Clitoria ternatea* L. *Taxonomic Serial No.:* 26543. https://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=26543#null.
- Aera, K., Jisoo, L., Nam, H. and Gyu Lee, H. 2016. Stabilization of black soybean anthocyanin by chitosan nanoencapsulation and copigmentation. *Journal of Food Biochemistry*, 41(2): 1-8. DOI: 10.1111/jfbc.12316.
- Alappat, B. and Alappat, J. 2020. Anthocyanin pigments: beyond aesthetics. *Molecules Journal*, 25: 1-17. <http://dx.doi.org/10.3390/molecules25235500>.
- Amanda, A. and Kurniaty, I. 2017. Pengaruh waktu maserasi terhadap rendemen zat antosianin pewarna alami minuman jelly dari terong ungu: *Seminar Nasional Sains dan Teknologi*, 19(12): 1-7. ISSN 2460-8416.
- Amaolo, A. 2017. The butterfly pea flower as a pH indicator. *International Scholastic Journal of Science*, 11(1): 1-3.
- Amperawati, S., Hastuti, P., Pranoto, Y. and Santoso, U. 2018. The anthocyanins content, colour changes and thermal stability of roselle (*Hibiscus sabdariffa* L.) petal extract. *International Journal of Science and Research*, 8(4): 428-435. DOI: 10.21275/ART20196610.
- Angriani, L. 2019. Potensi ekstrak bunga telang (*Clitoria ternatea*) sebagai pewarna alami local pada berbagai industry pangan. *Canrea Journal*, 2(2): 32-37.
- Barani, Y.H., Bhandari, B., Zhang, M. and Wang, B. 2020. Color stability and anthocyanins retention in microwave-thermally treated rose powder extracts during storage. *Journal of Food Processing and Preservation*, 44(10): 1-19. <http://dx.doi.org/10.1111/jfpp.14727>.
- Baskaran, A., Mudalib, S.K.A. and Izirwan, I. 2019. Optimization of aqueous extraction of blue dye from butterfly pea flower. *Journal of Physics*, 1358: 1-8. DOI: 10.1088/1742-6596/1358/1/012001.

Cai, L. 2014. Thin Layer Chromatography: *Current Protocols Essential Laboratory Techniques*. USA: Wiley Online Library.
<http://dx.doi.org/10.1002/9780470089941.et0603s08>.

Charurungsipong, P., Tangduangdee, C., Amornraksa, S., Asavasanti, S. and Lin J. 2020. Improvement of anthocyanin stability in butterfly pea flower extract by co-pigmentation with catechin. *E3S Web of Conferences*, 141: 1-4.
<https://doi.org/10.1051/e3sconf/202014103008>.

Chen, L., Chen I., Chen, P.Y., Huang, P. 2018. Application of butterfly pea flower extract in mask development. *Scientia Pharmaceutica*, 86(4): 1-9.
<http://dx.doi.org/10.3390/scipharm86040053>.

El-banna, F.S., Mahfouz, M.E., Loporatti, S., El-kemary, M. and Hanafy, N.A.N. 2019. Chitosan as a natural copolymer with unique properties for the development of hydrogels. *Appl. Sci.*, 9(11): 2193.
<https://doi.org/10.3390/app9112193>.

Fernandes, R.P.P., Trindade, M.A., Tonin, F.G., Lima, C.G., Pugine, S.M.P., Munekata, P.E.S., Lorenzo, J.M. and Melo, M.P. 2016. Evaluation of antioxidant capacity of 13 plant extracts by three different methods: cluster analyses applied for selection of the natural extracts with higher antioxidant capacity to replace synthetic antioxidant in lamb burgers. *J Food Sci Technol.*, 53(1): 451-460.

Fitriyani, R., Lestario, L.N. and Martono, Y. 2018. Jenis dan kandungan antosianin buah tomi-tomi. *J. Teknol. dan Industri Pangan*, 29(2): 137-144. DOI: 10.6066/jtip.2018.29.2.137.

Ge, J., Yue, P., Chi, J., Liang, J. and Gao, X. 2017. Formation and stability of anthocyanins-loaded nanocomplexes prepared with chitosan hydrochloride and carboxymethyl chitosan. *Food Hydrocolloids*, 74: 24-31.
<https://doi.org/10.1016/j.foodhyd.2017.07.029>.

Gençdag, E., Ozdemir, E.E., Demirci, K., Gorguc, A. and Yilmaz, F.M. 2022. Copigmentation and stabilization of anthocyanins using organic molecules and encapsulation techniques. *Current Plant Biology*, 29: 1-11.
<https://doi.org/10.1016/j.cpb.2022.100238>.

Goh, S.E., Kwong, P.J., Ng, C.L., Ng, W.J. and EE. K.Y. 2022. Antioxidant-rich *Clitoria ternatea* L. flower and its benefits in improving murine reproductive performance. *Food Sci. Technol.*, 42: 1-7.

Gonzalo, M. 2020. Copigmentation with chlorogenic and ferulic acid affected color and anthocyanin stability in model beverages colored with *Sambucus*

- peruviana*, *Sambucus nigra*, and *Daucus carota* during storage. *Foods*, 9(10): 1476. <http://dx.doi.org/10.3390/foods9101476>.
- Gordillo, B., Rodriguez-Pulido, F.J., Escudero-Gilete, M.L., Gonzalez-Miret, M.L. and Heredia, F.J. 2012. Comprehensive colorimetric study of anthocyanic copigmentation in model solutions. effects of ph and molar ratio. *Journal of Agricultural and Food Chemistry*, 60(11): 2896–2905. <http://dx.doi.org/10.1021/jf2046202>.
- Hariadi, H., Karim, M.A., Hanifah, U., Haryanto, A., Novrinaldi, Surahman, D.N., Astro, H.M., Assalam, S. and Lubis, R.F.L. 2022. Effect of butterfly-pea powder (*Clitoria ternatea* L.) and drying temperature towards physicochemical characteristics of butterfly-pea milk powder with vacuum drying method. *Food Sci. Technol.*, 43: 1-8. <https://doi.org/10.1590/fst.109622>.
- He, X., Li, X., LV, Y., and He, Q. 2015. Composition and color stability of anthocyanin-based extract from purple sweet potato: *Food. Sci. Technol.*, 35(3): 468-473. <http://dx.doi.org/10.1590/1678-457X.6687>.
- Husna, A., Lubis, Y.M. and Erika, C. 2022. Ekstraksi pewarna alami dari bunga telang (*Clitoria ternatea* L.) Dengan variasi jenis pelarut dan lama ekstraksi. *Jurnal Ilmiah Mahasiswa Pertanian*, 7(2): 410-418.
- Jadhav, R.V. and Bhujbal, S.S. 2020. A review on natural food colors. *Pharmaceutical Resonance*, 2(2): 12-20.
- Jeyaraj, E.J., Lim, Y.Y. and Choo, W.S. 2021. Extraction methods of butterfly pea (*Clitoria ternatea*) flower and biological activities of its phytochemicals. *J Food Sci Technol.*, 58(6): 2054-2067. <https://doi.org/10.1007%2Fs13197-020-04745-3>.
- Jung, H., Lee, Y., and Yoon, W. 2018. Effect of moisture content on the grinding process and powder properties in food: a review. *Processes*, 6(69): 1-16. <http://dx.doi.org/10.3390/pr6060069>.
- Kang, H.J., Ko, M.J. and Chung, M.S. 2021. Anthocyanin structure and pH dependent extraction characteristics from blueberries (*Vaccinium corymbosum*) and chokeberries (*Aronia melanocarpa*) in subcritical water state. *Foods*, 10(3): 1-12. <http://dx.doi.org/10.3390/foods10030527>.
- Khoo, H.E., Azlan, A., Tang, S.T. and Lim, S.M. 2017. Anthocyanidins and anthocyanins: colored pigments as food, pharmaceutical ingredients, and the potential health benefits. *Food and Nutrition Research*, 61: 1-21. <https://doi.org/10.1080/16546628.2017.1361779>.

- Kim, M.K., Kim, H.A., Koh, K. and Kim, H. 2018. Identification and quantification of anthocyanin pigments in colored rice. *Nutrition Research and Practice*, 2(1): 46-49. <http://dx.doi.org/10.4162/nrp.2008.2.1.46>.
- Ko, A., Lee, J.S., Nam, H.S. and Lee, H.G. 2016. Stabilization of black soybean anthocyanin by chitosan nanoencapsulation and copigmentation. *Journal of Food Biochemistry*, 41(2): 1-8. DOI: 10.1111/jfbc.12316.
- Kuppuswami, G.M. 2014. Fermentation (Industrial) – Production of Xanthan Gum: *Encyclopedia of Food Microbiology (Second Edition)*, pp. 816-821. <https://doi.org/10.1016/B978-0-12-384730-0.00110-5>.
- Lv, X., Mu, J., Wang, W., Liu, Y., Lu, X., Sun, J., Wang, J. and Ma, Q. 2022. Effects and mechanism of natural phenolic acids/fatty acids on copigmentation of purple sweet potato anthocyanins. *Current Research in Food Science*, 5: 1243-1250. <https://doi.org/10.1016%2Fcrfs.2022.08.003>.
- Marpaung, A.M., Lee, M. and Kartawiria, I.S. 2020. The development of butterfly pea (*clitoria ternatea*) flower powder drink by co-crystallization. *Indonesian Food Science and Technology Journal*, 3(2): 34-37. <https://doi.org/10.22437/ifstj.v3i2.10185>.
- Malacara, D. 2011. Color Vision and Colorimetry: Theory and Applications, Second Edition. SPIE Press: Bellingham, WA.
- Martins, N., Roriz, C.L., Mroales, P., Barros, L. and Ferreira, I.C.F.R. 2016. Food colorants: challenges, opportunities and current desires of agro-industries to ensure consumer expectations and regulatory practices: *Trends in Food Science and Technology*, 16: 1-25. <http://dx.doi.org/10.1016/j.tifs.2016.03.009>.
- Muthu, M., Gopal, J., Chun, S., Davadoss, A.J.P., Hasan, N. and Sivanasen, I. 2021. Crustacean waste-derived chitosan: antioxidant properties and perspectives. *Antioxidants*, 10(2): 228. <https://doi.org/10.3390/antiox10020228>.
- Nassour, R., Al-tameemi, K. and Ayash, A. 2020. Anthocyanin pigments: Structure and biological importance. *Journal of Chemical and Pharmaceutical Sciences*, 13(4): 45-57. ISSN 0974-2115.
- Oguis, G.K., Gilding, E.K., Jackson, M.A. and Craik, D.J. 2019. Butterfly pea (*Clitoria ternatea*), a cyclotide-bearing plant with applications in agriculture and medicine: *Front. Plant Sci.*, 10(645): 1-23. <https://doi.org/10.3389/fpls.2019.00645>.
- Oliveira, L.M. and Antelo, F. 2020. Thermostability of the visual color and anthocyanins from Rio-Grande-Cherry (*Eugenia involucrata* DC). *Brazilian*

Journal of Food Technology, 23: 1-13. <https://doi.org/10.1590/1981-6723.14019>.

Pertiwi, R.B., Hasbullah, U.H.A. and Affandi, A.R. 2021. Copigmentation of anthocyanin extract from parijoto fruit (*Medinilla speciosa*) and its stability at different temperatures and heating durations. *Indonesian Food and Nutrition Progress*, 18(2): 50-59. DOI: 10.22146/ifnp.65771.

Phrueksanan, W., Yibchok-anun, S. and Adisakwattana, S. 2014. Protection of *Clitoria ternatea* flower petal extract against free radical-induced hemolysis and oxidative damage in canine erythrocytes. *Res. Vet. Sci.*, 97: 357-363. DOI: 10.1016/j.rvsc.2014.08.010.

Picariello, L., Errichiello, F., Coppola, F., Rinaldi, A., Moio, L. and Gambuti, A. 2021. Effect of chitosan on the removal of different types of tannins from red wines: *Journal of Applied Science*, 11: 1-16. <https://doi.org/10.3390/app112411743>.

Pinheiro, C.P., Moreira, L.M.K., Alves, S.S., Jr, T.R.S.C. and Pinto, L.A.A. 2021. Anthocyanins concentration by adsorption onto chitosan and alginate beads: isotherms, kinetics and thermodynamics parameters. *International Journal of Biology Macromolecules*, 166: 934-939. <https://doi.org/10.1016/j.ijbiomac.2020.10.250>.

Rakić, V. and Ulrih, N.P. 2021. Influence of pH on color variation and stability of cyanidin and cyanidin 3-O- β -glucopyranoside in aqueous solution. *Journal of Food*, 19(1): 174-182. <https://doi.org/10.1080/19476337.2021.1874539>.

Ramadan, K.M.A. and El-Hadidy, E.M. 2015. Color stability of anthocyanin-based extracts in non-traditional sources: improvement of thermal stability by tannic acid. *J. Biol. Chem. Environ. Sci.*, 10(3): 1-19.

Ramadhani, R., Putra, G.G.P. and Suhendra, L. 2019. Karakteristik ekstrak kulit biji kakao (*Theobroma cacao* L.) sebagai sumber antioksidan pada perlakuan ukuran partikel dan waktu maserasi. *Jurnal Rekayasa dan Manajemen Argoindustri*, 9(2): 246-256.

Rodriguez-Amaya, D.B. and Carle, R. 2021. Chapter 7 – Alterations of natural pigments: Chemical Changes During Processing and Storage of Foods, pp. 265-327. USA: Academic Press. <https://doi.org/10.1016/B978-0-12-817380-0.00007-5>.

Sampebarra, A.L. 2018. Karakteristik zat warna antosianin dari biji kakao non fermentasi sebagai sumber zat warna alam. *Jurnal Industri Hasil Perkebunan*, 13(1): 63-70.

- Sánchez-Machado, D.I., López-Cervantes, J., Correa-Murrieta, M.A., Sánchez-Duarte, R.G., Cruz-Flores, P. and Mora-López, G.S. 2019. Chapter 4.2 – Chitosan: *Nonvitamin and Nonmineral Nutritional Supplements*, pp. 485-493. USA: Academic Press. <https://doi.org/10.1016/B978-0-12-812491-8.00064-3>.
- Santiago, M. and Strobel, S. 2013. Chapter 24 – Thin Layer Chromatography: *Methods in Enzymology*, pp. 303-324. USA: Academic Press. <https://doi.org/10.1016/B978-0-12-420067-8.00024-6>.
- Saptarini, N., Suryasaputra, D. and Nurmalia, H. 2015. Application of butterfly pea extract (*Clitoria ternatea* Linn) as an indicator of acid-base titration. *Journal of Chemical and Pharmaceutical Research*, 7(2): 275-280.
- Sharma, A., Gautam, S. and Wadhawan, S. 2014. Xanthomonas: *Encyclopedia of Food Microbiology (Second Edition)*, pp. 811-817. USA: Academic Press.
- Silva, V., Singh, R.K., Gomes, N., Soares, B.G., Silva, A., Falco, V., Capita, R., Alonso-Calleja, C., Pereira, J.E., Amaral, J.S., Igrelas, G. and Poeta, P. 2020. comparative insight upon chitosan solution and chitosan nanoparticles application on the phenolic content, antioxidant and antimicrobial activities of individual grape components of sousão variety. *Antioxidant*, 9(178): 1-14. <http://dx.doi.org/10.3390/antiox9020178>.
- Singh, M.V.P., Sanjuvikasini, V., Shruthi, S., Rajamehala, M., and Chozhavendhan, S. 2022. Chapter 23 – Value-added seafood products processing through novel enzyme: *Value-Addition in Food Products and Processing Through Enzyme Technology*, pp. 309-320. USA: Academic Press. <https://doi.org/10.1016/B978-0-323-89929-1.00013-5>.
- Singhvi, G., Hans, N., Shiva, N. and Dubey, S.K. 2019. Chapter 5 – Xanthan gum in drug delivery applications: *Natural Polysaccharides in Drug Delivery and Biomedical Application*, pp. 121-144. USA: Academic Press. <https://doi.org/10.1016/B978-0-12-817055-7.00005-4>.
- Suarna, I.W. and Wijaya, I.M.S. 2021. Butterfly pea (*Clitoria ternatea* L.: Fabaceae) and its morphological variations in Bali. *Journal of Tropical Biodiversity and Biotechnology*, 6(2): 63013. DOI: <http://dx.doi.org/10.22146/jtbb.63013>.
- Sugiarto, B., Rirung, Y., Ardiyani, A.N. and Achmad, Z. 2022. Extraction of natural color of butterfly pea (*Clitoria ternatea* L.) with variables pH-temperature and concentration for food coloring. *Journal Techno*, 8(1): 75-82.
- Sui, X., Dong, X. and Zhou, W. 2014. Combined effect of pH and high temperature on the stability and antioxidant capacity of two anthocyanins in aqueous

- solution. *Food Chemistry*, pp. 163-170. <http://dx.doi.org/10.1016/j.foodchem.2014.04.075>.
- Sultana, B., Anwar, F. and Ashrap, M. 2019. Effect of extraction solvent/technique on the antioxidant activity of selected medicinal plant extracts: *Molecules*, 14(6): 2167-2180. <https://doi.org/10.3390%2Fmolecules14062167>.
- Surianti, Husain, H. and Sulfikar. 2019. Uji stabilitas pigmen merah antosianin dari daun jati muda (*Tectona grandis* Linn f) terhadap pH sebagai pewarna alami. *Jurnal Chemica*, 20(1): 94-101.
- Suzery, M., Nudin, B., Bima, D.N., Cahyono, B. 2020. Effects of temperature and heating time on degradation and antioxidant activity of anthocyanin from roselle petals (*Hibiscus sabdariffa* L.). *International Journal of Science, Technology & Management*, 10: 289-296. ISSN:2722-4015.
- Teixeria-Costa, B. and Andrade, C.T. 2021. Chitosan as a valuable biomolecule from seafood industry waste in the design of green food packaging: *Biomolecules*, 11(11): 1599. <http://dx.doi.org/10.3390/biom11111599>.
- Thuy, M.N., Ben, T.C., Minh, V.Q. and Ngo, T.V. 2021. Identification of anthocyanin compounds in butterfly pea flowers (*Clitoria ternatea* L.) by ultra performance liquid chromatography/ultraviolet coupled to mass spectrometry. *Molecules*, 26(15): 1-13. <https://doi.org/10.3390/molecules26154539>.
- Unawahi, S., Widyasanti, A. and Rahimah, S. 2022. Pemanfaatan ekstrak bunga telang (*Clitoria ternatea* Linn) sebagai pewarna alami pada minuman bersoda. *Agrointek*, 16(2): 256-263. DOI 10.21107/agrointek.v16i2.13033.
- Utami, Y.P., Umar, A.H., Syahruni, R. and Kadullah, I. 2017. Standarisasi simplia dan ekstrak etanol daun leilem (*Clerodendrum minahassae* Teijsm. & Binn.). *Journal of Pharmaceutical and Medicinal Science*, 2(1): 32-39.
- Wahyuni, H., Hanum, T., and Murhani. 2017. Pengaruh kopigmentasi terhadap stabilitas warna antosianin ekstrak kulit terung belanda (*Cyphomandra betacea* Sendtn). *Jurnal Teknologi Industri dan Hasil Pertanian*, 22(1): 40-51.
- Wang, A. and Wang, W. 2013. Gum-g-Copolymers: Synthesis, Properties, and Applications: *Polysaccharide Based Graft Copolymers*, pp. 149-203. USA: Springer. http://dx.doi.org/10.1007/978-3-642-36566-9_5.
- Widelska, G. 2019. Impact of xanthan gum addition on phenolic acids composition and selected properties of new gluten-free maize-field bean pasta. *Open Chem.*, 17: 587-598. <https://doi.org/10.1515/chem-2019-0075>.

Wiyantoko, B. and Astuti. 2020. Butterfly pea (*Clitoria ternatea* L.) extract as indicator of acid-base titration. *Ind. J. Chem. Anal.*, 3(1): 22-32. ISSN 2622-7401.

Zhao, L., Pan, F., Mehmood, A., Zhang, Y., Hao, S., Ur, A., Li, J., Wang, C. and Wang, Y. 2020. Protective effect and mechanism of action of xanthan gum on the color stability of black rice anthocyanins in model beverage systems. *International Journal of Biology Macromolecules*, 164: 3800-3807. <https://doi.org/10.1016/j.ijbiomac.2020.09.027>.

Zhao, L., Pan, F., Mehmood, A., Zhang, H., Rehman, A., Li, J., Hao, S. and Wang, C. 2021. Improved color stability of anthocyanins in the presence of ascorbic acid with the combination of rosmarinic acid and xanthan gum. *Food Chemistry*, 35: 1-9. <https://doi.org/10.1016/j.foodchem.2021.129317>.

Zhu, Y., Chen, H., Lou, L., Chen, Y., Ye, X. and Chen, J. 2020. Copigmentation effect of three phenolic acids on color and thermal stability of Chinese bayberry anthocyanins. *Food Science Nutrition Journal*, 8(7): 3234-3242. <https://doi.org/10.1002%2Ffsn3.1583>.

Zingare, M.L., Zingare, P.L., Dubey, A.K. and Ansari, M.A. 2013. *Clitoria ternatea*: a review of the antioxidant, antidiabetic and hepatoprotective potentials, 3(1): 203-213. ISSN 2230-7605.