

DAFTAR PUSTAKA

1. Jilma B, Blann A, Pernerstorfer T, Stohlawetz P, Eichler H, Vondrovec B, et al. Human Endotoxemia. *Respir Critical Care Med*. 1999;1(14):857–63.
2. Zahorec R. Ratio of neutrophil to lymphocyte counts--rapid and simple parameter of systemic inflammation and stress in critically ill. *Bratisl Lek Listy [Internet]*. 2001 [cited 2017 Sep 21];102(1):5–14. Available from: <http://bmj.fmed.uniba.sk/2001/10201-01.PDF>
3. Lowsby R, Gomes C, Jarman I, Lisboa P, Nee PA, Vardhan M, et al. Neutrophil to lymphocyte count ratio as an early indicator of blood stream infection in the emergency department. *Emerg Med J [Internet]*. 2015;32(7):531–4. Available from: <http://emj.bmj.com/lookup/doi/10.1136/emered-2014-204071>
4. Li H, Lu X, Xiong R, Wang S. High Neutrophil-to-Lymphocyte Ratio Predicts Cardiovascular Mortality in Chronic Hemodialysis Patients. 2017;2017(Lvmi).
5. Giede-Jeppe A, Bobinger T, Gerner ST, Sembill JA, Sprügel MI, Beuscher VD, et al. Neutrophil-to-Lymphocyte Ratio Is an Independent Predictor for In-Hospital Mortality in Spontaneous Intracerebral Hemorrhage. *Cerebrovasc Dis*. 2017;44(1–2):26–34.
6. de Jager CPC, Wever PC, Gemen EFA, Kusters R, van Gageldonk-Lafeber AB, van der Poll T, et al. The Neutrophil-Lymphocyte Count Ratio in Patients with Community-Acquired Pneumonia. *PLoS One*. 2012;7(10).
7. Saliccioli JD, Marshall DC, Pimentel M, Santos MD, Pollard T, Celi L, et al. The association between the neutrophil-to-lymphocyte ratio and mortality in critical illness: an observational cohort study. *Crit Care [Internet]*. 2015;19(1):13. Available from: <http://ccforum.com/content/19/1/13>
8. Adi Nugroho, Suwarman AMN. Hubungan antara Rasio Neutrofil-Limfosit dan Skor Sequential Organ Failure Assesment pada Pasien yang Dirawat di Ruang Intensive Care Unit. *Anestesi Perioper*. 2013;1(3):189–96.
9. Abbas AK, Lichtman AH, Pillai S. *Cellular and Molecular Immunology*. 8th ed. Canada: Elsevier; 2014.
10. Kruger P, Saffarzadeh M, Weber ANR, Rieber N, Radsak M, von Bernuth H, et al. Neutrophils: Between Host Defence, Immune Modulation, and Tissue Injury. *PLoS Pathog*. 2015;11(3):1–22.

11. Dionigi R, Dominioni L, Benevento A, Giudice G, Cuffari S, Bordone N, et al. Effects of surgical trauma of laparoscopic vs. open cholecystectomy. *Hepatogastroenterology* [Internet]. 1994 Oct;41(5):471—476. Available from: <http://europepmc.org/abstract/MED/7851857>
12. Ayala A, Herdon CD, Lehman DL, Ayala C a, Chaudry IH. Differential induction of apoptosis in lymphoid tissues during sepsis: variation in onset, frequency, and the nature of the mediators. *Blood*. 1996;87(10):4261–75.
13. Menges T, Engel J, Welters I, Wagner R-M, Little S, Ruwoldt R, et al. Changes in blood lymphocyte populations after multiple trauma: Association with posttraumatic complications. *Crit Care Med* [Internet]. 1999;27(4). Available from: http://journals.lww.com/ccmjournal/Fulltext/1999/04000/Changes_in_blood_lymphocyte_populations_after.26.aspx
14. Sterzl J. Vývoj a indukce imunitní odpovědi. Praha: Academia; 1988.
15. Wang X, Zhang G, Jiang X, Zhu H, Lu Z, Xu L. Neutrophil to lymphocyte ratio in relation to risk of all-cause mortality and cardiovascular events among patients undergoing angiography or cardiac revascularization: A meta-analysis of observational studies. *Atherosclerosis* [Internet]. 2014;234(1):206–13. Available from: <http://www.sciencedirect.com/science/article/pii/S0021915014001488>
16. Huehnergath K V, Mozaffarian D, Sullivan MD, Crane BA, Wilkinson CW, Lawler RL, et al. Usefulness of Relative Lymphocyte Count as an Independent Predictor of Death/Urgent Transplant in Heart Failure. *Am J Cardiol* [Internet]. 2005;95(12):1492–5. Available from: <http://www.sciencedirect.com/science/article/pii/S0002914905004820>
17. Papa A, Emdin M, Passino C, Michelassi C, Battaglia D, Cocci F. Predictive value of elevated neutrophil-lymphocyte ratio on cardiac mortality in patients with stable coronary artery disease. *Clin Chim Acta* [Internet]. 2008 [cited 2017 Sep 21];395(1–2):27–31. Available from: <http://www.sciencedirect.com/science/article/pii/S0009898108002143>
18. Walsh SR, Cook EJ, Goulder F, Justin TA, Keeling NJ. Neutrophil-lymphocyte ratio as a prognostic factor in colorectal cancer. *J Surg Oncol* [Internet]. 2005 Sep 1 [cited 2017 Sep 21];91(3):181–4. Available from: <http://doi.wiley.com/10.1002/jso.20329>
19. Tamhane U, Aneja S, Montgomery D. Association between admission neutrophil to lymphocyte ratio and outcomes in patients with acute coronary

syndrome. *Am J* [Internet]. 2008 [cited 2017 Sep 21]; Available from: <http://www.sciencedirect.com/science/article/pii/S0002914908008266>

20. Neuen BL, Leather N, Greenwood AM, Gunnarsson R, Cho Y, Mantha ML. Neutrophil–lymphocyte ratio predicts cardiovascular and all-cause mortality in hemodialysis patients. *Ren Fail* [Internet]. 2016;38(1):70–6. Available from: <http://www.tandfonline.com/doi/full/10.3109/0886022X.2015.1104990>

