

DAFTAR PUSTAKA

1. Smeltzer SCO, Bare BG, Hinkle JL, Cheever KH. Brunner & Suddarth's Textbook of Medical-surgical Nursing, Volume 1, 12th ed.: Lippincott Williams & Wilkins; 2010.
2. Sheta SA. Procedural sedation analgesia. Saudi Journal of Anaesthesia. 2010;4(1):11-16.
3. Barash PG, Cullen BF, Stoelting RK, Cahalan M, Stock MC. Clinical Anesthesia, 6th ed. Philadelphia: Lippincott Williams & Wilkins; 2009.
4. Poh CF, Ng S, Berean KW, Williams PM, Rosin MP, Zhang L. Biopsy and histopathologic diagnosis of oral premalignant and malignant lesions. J Can Dent Assoc. 2008;74:283–288.
5. Logan RM, Goss AN. Biopsy of the oral mucosa and use of histopathology services. Aust Dent J. 2010;55:9–13.
6. Caddy C, Giaroli G, White TP, Shergill SS, Tracy DK. Ketamine as the prototype glutamatergic antidepressant: pharmacodynamic actions, and a systematic review and meta-analysis of efficacy. Therapeutic Advances in Psychopharmacology. 2013;4(2):75-99.
7. Brown EN, Lydic R, Schiff ND. General Anesthesia, Sleep, and Coma. N Engl J Med. 2010;363:2638-50
8. Andropoulos DB, Greene MF. The Anesthesia and Developing Brains — Implications of the FDA Warning; N Engl J Med. 2017;376(1):1-3.
9. Kim KM, Lee KH, Kim YH, Ko MJ, Jung JW, Kang E. Comparison of effects of intravenous midazolam and ketamine on emergence agitation in children: Randomized controlled trial. Journal of International Medical Research 2016;44(2):258-266.
10. Lonardo NW, Mone MC, Nirula R, Kimball EJ, Ludwig K, Zhou X, Sauer BC, Nechodom K, Teng C, Barton RG. Propofol Is Associated with Favorable Outcomes Compared with Benzodiazepines in Ventilated Intensive Care Unit Patients. 2014;189(11):1383-1394.

11. Ghezzi F, Monni L, Corsini S, Rauti R, Nistri A. Propofol protects rat hypoglossal motoneurons in an in vitro model of excitotoxicity by boosting GABAergic inhibition and reducing oxidative stress, *Neuroscience*. 2017;367:15-33.
12. Krithika V, Amudharani R, Anandan H. Comparison of Ketamine and Propofol in Combination with keta and Midazolam in Total Intravenous Anesthesia for Minor Gynecological Procedures. *Int J Sci Stud.* 2017;5(2):178-180.
13. Chan S-M, Lee M-S, Lu C-H, Cherng C-H, Huang Y-S, Yeh C-C, et al. Confounding Factors to Predict the Awakening Effect-Site Concentration of Propofol in Target-Controlled Infusion Based on Propofol and Fentanyl Anesthesia. *PLoS ONE*. 2015;10(5):e0124343.
14. Deschka M, Trein TA, Wagatsuma JT, Silva GL, Araújo MA, Games BMM, Alcindo JF, Celso Antônio Rodrigues CA, Santos PSP. Bispectral index assessment in calves subjected to the continuous infusion of propofol combined with fentanyl administration. 2017; 47(11):e20151621.
15. Farahmand F, Roze ME, Gudarzi M, Najafi M, Fallahi G, Motamed F, et al. Evaluation of the Effect of Intravenous Ondansetron versus Placebo before Anesthesia on Vomiting after Endoscopy and Colonoscopy Procedures. *Acad J Surg*, 2017; 4(1): 6-8.
16. Pendi A, Field R, Farhan SD, Eichler M, Bederman SS. Perioperative ketamine for analgesia in spine surgery: a meta-analysis of randomized controlled trials. *Spine*. 2018;43:E299–E307.
17. Chen, J-Y., Jia, J-E., Liu, T-J. et al. *Can J Anesth/J Can Anesth* 2013; 60: 385.
18. Green SM, Roback MG, Kennedy RM, Krauss B. Clinical Practice Guideline For Emergency Department Ketamine Dissociative Sedation: 2011 Update. 2011;57:449–461.
19. Martin DC, Watkins CA, Adams RJ, Nason LA. Anesthetic effects on 5-hydroxytryptamine uptake by rat brain synaptosomes. *BrainRes*. 1988;455: 360–5

20. Tsai H-C, Lin Y-C, Ko C-L, Lou H-Y, Chen T-L, Tam K-W, et al. Propofol versus Midazolam for Upper Gastrointestinal Endoscopy in Cirrhotic Patients: A Meta-Analysis of Randomized Controlled Trials. *PLoS ONE*. 2015;10(2): e0117585.
21. Shaikh SI, Nagarekha D, Hegade G, Marutheesh M. Postoperative nausea and vomiting: A simple yet complex problem. *Anesthesia, Essays and Researches*. 2016;10(3):388-396.
22. Motamed F, Aminpour Y, Hashemian H, Soltani AE, Najafi M, Farahmand F. Midazolam–ketamine combination for moderate sedation in upper GI endoscopy. *J Pediatr Gastroenterol Nutr*. 2012; 54:422–426.
23. Gómez F, Phillips C, Soddu A, Boly M, Boveroux P, Vanhaudenhuyse A, et al. Changes in Effective Connectivity by Propofol Sedation. *PLoS ONE*, 2013;8(8):e71370.
24. White PF, Song D-J. New Criteria for Fast-Tracking After Outpatient Anesthesia: A Comparison with the Modified Aldrete's Scoring System. 1999;88:1069–72
25. Aldrete JA. The post- anesthesia recovery score revisited. *J Clin Anesth*. 1995;7:89- 91.
26. Bhavsar S, Dang AQ. Anaesthetic considerations in measuring Aldrete score and long-term outcomes in craniotomy patients. *European Journal of Anaesthesiology (EJA)*. 2013; 30(9):576–577.
27. Lotia S, Bellamy MC. Anaesthesia and morbid obesity, Continuing Education in Anaesthesia Critical Care & Pain. 2008;8(5):151–156.
28. Kanonidou Z, Karytianou G. Anesthesia for the elderly. *Hippokratia*. 2007;11(4):175-177.
29. Mawe GM, Hoffman JM. Serotonin Signaling in the Gastrointestinal Tract: Functions, Dysfunctions, and Therapeutic Targets. *Nature Reviews Gastroenterology & Hepatology*. 2013;10(8):473-486.