

BIBLIOGRAPHY

- Adebiyi, A. P., Adebiyi, A. O., Yamashita, J., Ogawa, T., and Muramoto, K. 2008. Purification and characterization of antioxidative peptides derived from rice bran protein hydrolysates. European Food Research Technology 228: 553-563.
- Almeida, C. C., Monteiro, M.L.G., da Costa-Lima, B. R. C., Alvares, T. S., and Conte-Junior, C. A. 2014. In vitro digestibility of commercial whey protein supplements. Food Science and Technology 61(1): 7-11.
- AOAC. 2005. "Official Methods of Analysis of the Association of Official Analytical Chemists." AOAC Inc., Arlington.
- Badan Pusat Statistik. 2015. Laporan Statistik Pertanian Tanaman Pangan. Available from: <https://www.bps.go.id/linkTableDinamis/view/id/871>. Accessed 2018 May 5th.
- Bartosz, G. 2014. "Food oxidants and antioxidants: Chemical, biological, and functional properties". CRC Press, Boca Raton.
- Beuchat, L. R., Komitopoulou, E., Beckers, H., Betts, R. P., Bourdichon, F., Fanning, S., Joosten, H. M., and Kuile, B. H. T. Low-Water Activity Foods: Increased Concern as Vehicles of Foodborne Pathogens.
- Brown, A. 2010. "Understanding Food Principles & Preparation". Wadsworth, Belmont.
- Castro, R.J.S.D, and Sato, H.H. 2013. Advantages of an acid protease from *Aspergillus oryzae* over commercial preparations for production of whey protein hydrolysates with antioxidant activities. Biocatalysis and Agricultural Biotechnology 3(3): 58-65.
- Chen, H. 2013. "Modern Solid State Fermentation Theory and Practice". Springer, Dordrecht.
- Chen, L., Madi, R. L., and Vadlani, P. V. 2013. "Value-Added Products from Soybean: Removal of Anti-Nutritional Factors via Bioprocessing". InTech, Rijeka.
- Cheng, J. 2017. "Biomass to Renewable Energy Processes" 2nd ed. CRC Press, Boca Raton.
- Chutmanop, J., Chuichulcherm, S., Chisti, Y., and Srinophakun, P. 2008. Protease production by *Aspergillus oryzae* in solid-state fermentation using

- agroindustrial substrates. *Journal of Chemical Technology and Biotechnology* 83: 1012-1018.
- Cirilo, G., and Iemma, F. 2012. Antioxidant polymers: synthesis, properties, and applications. John Wiley & Sons, Beverly.
- Dharmawanti, R. K. W. A. 2009. Analisis Strategi Pengembangan Usaha Pada Primer Koperasi Produsen Tempe Tahu Indonesia. *Strata I* thesis, Univ. Pertanian Bogor.
- El-Fadaly, H. M., El-Kadi, S. M., Hamad, M. N., and Habib, A. A. 2015. Isolation and Identification of Egyptian Ras Cheese (Romy) Contaminating Fungi during Ripening Period. *Journal of Microbiology Research* 5(1): 1-10.
- He, R., Ju, X., Yuan, J., Wang, L., Girgih, A. T., and Aluko, R. E. 2012. Antioxidant activities of rapeseed peptides produced by solid state fermentation. *Food Research International* 49: 432-438.
- Hur, S. J., Lee, S.Y., Kim, Y-C., Choi, I., and Kim, G-B. 2014. Effect of fermentation on the antioxidant activity in plant-based foods. *Food Chemistry* 160: 346-56.
- Iyer, P., and Singhal, R. S. 2008. Production of glutaminase (E.C.3.2.1.5) from *Zygosaccharomyces rouxii*: Statistical optimization using response surface methodology. *Bioresource Technology* 99: 4300-4307.
- Jiménez-Escríg, A., Alaiz, M., Vioque, J., and Rupérez P. 2009. Health-promoting activities of ultra-filtered okara protein hydrolysates released by in vitro gastrointestinal digestion: identification of active peptide from soybean lipoxygenase. *European Food Research and Technology* 230(4): 655-63.
- Kim, S.K. 2012. "Handbook of Marine Macroalgae: Biotechnology and Applied Phycology". John Wiley & Sons, Ltd., Hoboken.
- Kuo, H. T., Wu, P. H., Chung, H. H., Jiang, C. M., and Wu, M. C. 2013. Koji fermentation improve the protein digestibility. *Journal of Food Processing and Preservation* 37(5): 419-423.
- Kwon, D. Y., Daily III, J. W., Kim, H. J., and Park, S. 2010. Antidiabetic effects of fermented soybean products on type 2 diabetes. *Nutrition research* 30: 1-13.
- Li, B., Qiao, M., and Lu, F. 2012. Composition, Nutrition, and Utilization of Okara (Soybean Residue). *Food Reviews International* 28(3): 231-52.
- Li, B., Zhang, Y., Yang, H., and Li, R. 2008. Effect of drying methods on functional properties of bean curd dregs. *J. Henan Inst. Science Technology* 36(3): 64-66.

- Lopez, O.P. 2017. "Amaranth Biology, Chemistry, and Technology". CRC Press. Boca Raton.
- Lu, F., Cui, Z., Liu, Yang., Li, Bo. 2013. The Effect of Okara on the Qualities of Noodle and Steamed Bread. Advance Journal of Food Science and Technology 5(7): 960-968.
- Manan, M. A., and Webb, C. 2017. Design Aspects of Solid State Fermentation as Applied to Microbial Bioprocessing. Journal of Applied Biotechnology & Bioengineering 4(1): 91-116.
- Mateos-Aparicio, I., Redondo-Cuenca A., Villanueva-Suárez M-J., Zapata-Revilla M-A., and Tenorio-Sanz M-D. 2010. Pea pod, broad bean pod and okara, potential sources of functional compounds. Food Science and Technology 43(9): 1467-470.
- McCarthy, A., Ocallaghan, Y., and Obrien, N. 2013. Protein Hydrolysates from Agricultural Crops—Bioactivity and Potential for Functional Food Development. Agriculture 3(1): 112-30.
- Membrillo, I., Sanchez, C., Meneses, M., Favela, E., and Loera, O. 2008. Effect of substrate particle size and additional nitrogen source on production of lignocellulolytic enzymes by *Pleurotus ostreatus* strains. Bioresource Technology 99: 7842-7847.
- Moayedi, A., Hashemi, M., and Safar, M. 2015. Valorization of tomato waste proteins through production of antioxidant and antibacterial hydrolysates by proteolytic *Bacillus subtilis*: optimization of fermentation conditions. Journal of Food Science and Technology 53(1): 391-400.
- Morais, H. A., Silvestre, M. P., Silveira, J. N., Silva, A. C., Silva, V. D., and Silva, M. R. 2013. Action of a pancreatin and an *Aspergillus oryzae* protease on whey proteins: Correlation among the methods of analysis of the enzymatic hydrolysates. Brazilian Archives of Biology and Technology 56(6): 985-995.
- Narayan, S. Meyers D. G., Graham K. S., Scott R., and Bennet M. J. 2011. Measurement of plasma amino acids by Ultraperformace[®] Liquid Chromatography. Clinical Chemistry and Laboratory Medicine. 49(7): 1177-1185.
- Oliveira, M. dos S., Feddern, V., Kupski, L., Cipolatti, E. P., Baditate-Furlong, E., and de Souza-Soares, L. A. 2010. Physico-chemical characterization of fermented rice bran biomass. CyTA Journal of Food 8(3): 229-236.
- Pangesti, N. W. I., Pangastuti, A., and Retnaningtyas, E. 2012. Pengaruh penambahan molase pada produksi enzim xylanase oleh fungi *Aspergillus niger* dengan substrat jerami padi. Bioteknologi 9 (2): 41-48.

Pasupuleti, V. K., and Demain, A.L. 2010. "Protein Hydrolysates in Biotechnology". Springer, New York.

Pownall, T. L., Udenigwe, C.C., and Aluko, R.E. 2010. Amino Acid Composition and Antioxidant Properties of Pea Seed (*Pisum sativum* L.) Enzymatic Protein Hydrolysate Fractions. *Journal of Agricultural and Food Chemistry* 58(8): 4712-718.

Rashad, M.M., Mahmoud, A.E., Abdou, H.M., and Nooman, M.U. 2011. Improvement of nutritional quality and antioxidant activities of yeast fermented soybean curd residue. *African Journal of Biotechnology* 10(28): 5504-513.

Salleh, A. B., Rahman, R.N.Z.R.A., and Basri, M. 2013. "Molecular and Structural Biology of New Lipases and Proteases". New York: Nova Science Publisher.

Sathya, R., Pradeep, B. V., Angaryakanni, J., and Palaniswamy, M. 2009. Production of Milk Clotting Protease by a Local Isolate of *Mucor circinelloides* under DDF using Agro-industrial Wastes. *Biotechnology and Bioprocess Engineering* 14: 788-794.

Sehu, K., and Bello, M. T. 2011. Effect of environmental factors on the growth of *Aspergillus species* associated with stored millet grains in sokoto. *Nigerian Journal of Basic and Applied Science* 19(2): 218-223.

Sengupta, S., Chakraborty, M., Bhowal, J., and Bhattacharya, D. K. 2012. Study on the effects of drying process on the composition and quality of wet okara. *International Journal of Science, Environment, and Technology* 1(4): 319-330.

Shahidi, F. 2015. "Handbook of Antioxidants for Food Preservation". Woodhead Publishing: Amsterdam.

Shurtleff, W., and Aoyagi, A. 2014. "Early History of Soybeans and Soyfoods Worldwide". Soyinfo, California.

Stirnemann, G., Kang, S. G., Zhou, R., and Berne B. J. 2014. How force unfolding differs from chemical denaturation. *Biophysics and Computational Biology, Chemistry* 111(9): 3413-3418.

Talebnia, F., Karakashev, D., and Angelidaki, I. 2010. Production of bioethanol from wheat straw: An overview on pretreatment, hydrolysis and fermentation. *Bioresource Technology* 1010: 4744-4753.

Teng, D., Gao, M., Yang, Y., Liu, B., Tian, Z., and Wang, J. 2012. Bio-modification of soybean meal with *Bacillus subtilis* or *Aspergillus oryzae*. *Biocatalysis and Agricultural Biotechnology* 1: 32-38.

USDA (United States Department of Agriculture). 2013. National Nutrient Database for Standard Reference Legacy Release (Okara, 16130). Available from:<https://ndb.nal.usda.gov/ndb/foods/show/16130?n1=%7BQv%3D1%7D&fgcd=&man=&lfacet=&count=&max=25&sort=default&qlookup=okara&offset=&format=Full&new=&measureby=&Qv=1&ds=&qt=&qp=&qa=&qn=&q=&ing=>. Accessed 2018 May 5th.

Vishwanathan, K. H., Govindaraju, K., Singh, V., and Subramanian, R. 2011. Production of okara and soy protein concentrates using membrane technology. Journal of Food Science 76(1): 158-164.

Vishwanatha, K. S., Rao, A. G. A., and Singh, S. A. 2010. Production and characterization of a milk-clotting enzyme from *Aspergillus oryzae* MTCC 5341. Applied Microbiology and Biotechnology 85(6): 1849-1859.

Yeh, A-I., Huang, Y-C., and Chen, S. H. 2010. Effect of particle size on the rate of enzymatic hydrolysis of cellulose. Carbohydrate Polymers 79: 192-199. \

Yoon, G., and Park, S. 2014. Antioxidant action of soy isoflavones on oxidative stress and antioxidant enzyme activities in exercised rats. Nutrition Research and Practice 8(6): 618-624.

Yu, K. W., Lee, S. E., Choi, H. S., Suh, H. J., Ra, K. S., Choi, J. W., and Hwang, J. H. 2012. Optimization for Rice *Koji* Preparation Using *Aspergillus oryzae* CJCM-4 Isolated from a Korean Traditional Meju. Food Science Biotechnology 21(1): 129-135.

Zhu, Y. P., Fan, J. F., Cheng, Y. Q., and Li, L. T. 2008. Improvement of the antioxidant activity of Chinese traditional fermented okara (Meitauza) using *Bacillus subtilis* B2. Food Control 19: 654-661.