

BIBLIOGRAPHY

- Aguilera, J. M., Simpson, R., Well-Chanes, J., Aguirre, D. B., and Barbosa-Cánovas, G. V. 2011. Food Engineering Interfaces. New York: Springer.
- Ahmad, A. F., and Youssef, M. S. H. 2015. Chemical composition and bioactive properties of *Illicium verum* (star-anise) extracts Prepared by different methods. Journal of Chem., Bio. and Phy. Sciences, 5(2); 1160–1170.
- Ahzar, Aina Afqah Binti. 2017. Optimization on Formula of Matcha Added Cajuputs Candy based on Total Phenolic Content and Sensory Acceptability. Thesis. Institut Pertanian Bogor.
- Al-Ghafari, A. B., Shorbaji, A. M., AL-Sarori, L. A., Baduwailan, E. O., Basaar, A. A., Al Doghaither, H. A., Omar, U. M. 2016. Phenolic Contents and Antioxidant Activities of Green Tea with and without Lemon. Natural Science, 8, 247–255.
- Ananingsih, V. K., Sharma, A., and Zhou, W. 2013. Green tea catechins during food processing and storage: A review on stability and detection. Food Res Int., 50(2); 469–479.
- Anesini C, Ferraro GE and Filip R. 2008. Total polyphenol content and antioxidant capacity of commercially available tea (*Camellia sinensis*) in Argentina. J Agr Food Chem 56: 9225–9229.
- AOAC. 2000. Official Methods of Analysis. 17th ed. Arlington: Association of Official Analytical Chemists.
- Ashardiono, F., and Cassim, M. 2015. Adapting to Climate Change: Challenges for Uji Tea Cultivation. J-Sustain, 3(1); 32–36.
- Astill, C., Birch, M. R., Dacombe, C., Humphrey, P. G. and Martin, P. T. 2001. Factors affecting the caffeine and polyphenol contents of black and green tea infusions. J. Agric. Food Chem, 49(11); 5340–7.
- Awad, O. I., Mamat, R., Ali, O. M., Azmi, W. H., Kadirkama, K., Yusri, I. M., Yusaf, T. 2017. Response surface methodology (RSM) based multi-objective optimization of fusel oil-gasoline blends at different water content in SI engine. Energy Convers. Manag, 150.
- Badan Standardisasi Nasional. 2008. Kembang Gula- Bagian 1: Keras. Standar Nasional Indonesia (SNI) 3547.1:2008. Jakarta: BSN.

- Bernal, M., Llorens, L., Julkunen-Tiltto, R., Badosa, J., and Verdaguer, D. 2013. Altitudinal and seasonal changes of phenolic compounds in *Buxus sempervirens* leaves and cuticles. *Plant Physiol Biochem*, 70, 471–482.
- Bizuayehu, D., Atlabachew, M. and Ali, M. T. 2016. Determination of some selected secondary metabolites and their in vitro antioxidant activity in commercially available Ethiopian tea (*Camellia sinensis*). *Springerplus*, 5, 412–421.
- Borse, B. B., Rao, L. J. M., Nagalakshmi, S. and Krishnamurthy, N. 2002. Fingerprint of black teas from India: Identification the regiospecific characteristics. *Food Chem.*, 79(4); 419–424.
- Bouayed, J., Hoffmann, L. and Bohn, T. 2011. Total phenolics, flavonoids, anthocyanins and antioxidant activity following simulated gastro-intestinal digestion and dialysis of apple varieties: Bioaccessibility and potential uptake. *Food Chem.*, 128(1); 14–21.
- Candioti, L. V., Zan, M. M. De, Camara, M. S. and Goicoechea, H. C. 2014. Experimental design and multiple response optimization. Using the desirability function in analytical methods development. *Talanta*, 124, 123–138.
- Cappa, C., Lavelli., V and Mariotti., M. 2015. Fruit candies enriched with grape skin powders: physicochemical properties. *LWT – Food Science and Tech.*, 62, 569 – 575.
- Carpenter, R. P., Lyon, D. H., and Hasdell, T. A. 2012. Guidelines for Sensory Analysis in Food Product Development and Quality Control (Second). Maryland: Aspen Publishers.
- Chacko, S. M., Thambi, P. T., Kuttan, R. and Nishigaki, I. 2010. Beneficial effects of green tea: A literature review. *Chinese Medicine*, 5(13); 1–9.
- Chatterjee, A., Saluja, M., Agarwal, G., and Alam, M. 2012. Green tea: A boon for periodontal and general health. *J. Indian Soc. Periodontol*, 16(2); 161–167.
- Chaturvedula, V. S. P. and Prakash, I. 2011. The aroma, taste, color and bioactive constituents of tea. *J. Med. Plants Res.*, 5, 2110–2124.
- Christie, G. 2012. Formulating on non sucrose flavoured “cajuput candy”. Thesis, Universitas Pelita Harapan, Tangerang.
- Deming, S. N. and Morgan, S. L. 2009. Experimental Design: A Chemometric Approach. Amsterdam: Elsevier Science Publishers.

- Fitrianto, A. and Midi, H. 2012. Multi-Response Optimization via Desirability Functionfor the Black Liquor Data. *J Sci Technol*, 4(1); 91–102.
- Gaylard, L. 2015. “The Tea Book”. New York: DK Publishing.
- Goran, M. I., Tappy, L. and Lê, K.-A. 2015. Dietary Sugars and Health. Boca Raton: CRC Press.
- Hartel, R. W., Elbe, J. H. von and Hofberger, R. 2017. Confectionery Science and Technology. Cham: Springer.
- Hayouni A., Bouix M., Abedrabba M., Leveau J. Y. and Hamdi M. 2008. Mechanism of action of *Melaleuca armillaris* (Sol. Ex Gaertn) Sm. essential oil on six LAB strains as assessed by multiparametric flow cytometry and automated microtiter-based assay. *Food Chem*. 111:707-718
- Horvathova, E., Turcaniova, V. and Slamenova, D. 2007. Comparative study of DNA-damaging and DNA-protective effects of selected components of essential plant oils in human leukemic cells K562. *Neoplasma*, 54(6); 478–83.
- Jeong, S.-M., Kim, S.-Y., Kim, D.-R., Jo, S.-C., Nam, K. C., Ahn, D. U. and Lee, S.-C. 2004. Effect of Heat Treatment on the Antioxidant Activity of Extracts from Citrus Peels. *J. Agric. Food Chem*, 52, 3389–3393.
- Jiang, B. and Zhang, Z.-W. 2012. Comparison on Phenolic Compounds and Antioxidant Properties of Cabernet Sauvignon and Merlot Wines from Four Wine Grape-Growing Regions in China. *Molecules*, 17, 8804–8821.
- Jothy, S. L., Zuraini, Z. and Sasidharan, S. 2011. Phytochemicals screening, DPPH free radical scavenging and xanthine oxidase inhibitory activities of *Cassia fistula* seeds extract. *J. Med. Plant. Res.*, 5(10); 1941–1947.
- Karori, S. M., Wachira, F. N., Wanyoko, J. K. and Ngure, R. M. 2007. Antioxidant capacity of different types of tea products. *African*, 6(19); 2287–2296.
- Kligler, B. and Chaudhary, S. 2007. Peppermint Oil. *Am Fam Physician*, 75, 1027–30.
- Kopjar, M., Tadić, M. and Piližota, V. 2015. Phenol content and antioxidant activity of green, yellow and black tea leaves. *Chemical and Biological Technologies in Agriculture*, 2, 1–6.
- Laaksonen, O. 2011. Astringent Food Compounds and Their Interactions with Taste Properties. Turku: University of Turku.

- Lambert, J. D. and Elias, R. J. 2010. The antioxidant and pro-oxidant activities of green tea polyphenols: a role in cancer prevention. *Arch Biochem Biophys*, 501(1); 65–72.
- Lawless, H. T. and Heymann, H. 2013. “Sensory Evaluation of Food: Principles and Practices”. New York: Springer.
- Lee, J., Chambers, D. H., Chambers IV, E., Adhikari, K. and Yoon, Y. 2013. Volatile Aroma Compounds in Various Brewed Green Teas. *Molecules*, 18, 10024–10042.
- Lorenzo, J.M. and Munekata, P.E.S. 2016. Phenolic compounds of green tea: Health benefits and technological application in food. *Asian Pac J Trop Biomed*, 6(8); 709-719.
- Mennella, J. A., Pepino, M. Y., Duke, F. F., & Reed, D. R. 2010. Age modifies the genotype-phenotype relationship for the bitter receptor TAS2R38. *BMC Genetics*, 11(60); 1–9.
- Mildner-Szkudlarz, S., Zawirska-Wojtasiak, R., Obuchowski, W. and Go, M. 2009. Evaluation of antioxidant activity of green tea extract and its effect on the biscuits lipid fraction oxidative stability. *JFS*, 74, 362-370.
- Moskowitz, H. R. 2006. “Applied Sensory Analysis of Foods”. Florida: CRC Press.
- Muzzaffar, S., Baba, Waqas, N., Nazir, N., Masoodi, F. A., Bhat, M. M., & Bazaz, R. 2016. Effect of storage on physicochemical, microbial and antioxidant properties of pumpkin (*Cucurbita moschata*) candy. *Cogent Food & Agriculture*, 2(1163650); 1–13.
- Myerson, A. 2002. “Handbook of Industrial Crystallization” 2nd ed. Woburn: Butterworth Heinemann.
- Norazalina. 2016. Improvement of Cajuputs Candy’s Acceptability by the Addition of Mango Flavouring and Natural Catechin Colourant. Thesis. Insitut Pertanian Bogor
- Pereira, V. P., Knor, F. J., Veloso, J. C. R., and Beltrame, F. L. 2014. Determination of phenolic compounds and antioxidant activity of green, black and white teas of *Camellia sinensis* (L.) Kuntze, *Theaceae*. *Revista Brasileira de Plantas Medicinais*, 16(3).
- Pusat Penelitian Teh dan Kina. 2014. Sejarah. Available from: <http://www.gamboeng.com/pages/detail/2014/29/109>. Accessed 2017 November 19.

Pusat Penelitian Teh dan Kina. 2015. Klon GMB 1-11. Available from: <http://www.gamboeng.com/pages/detail/2015/59/146>. Accessed 2017 December 8.

Raissi, S. and Farsani, R. E. 2009. Statistical Process Optimization Through Multi-Response Surface Methodology. *World Acad Sci Eng Technol*, 51, 267–271.

Rohdiana, D. 2015. Teh: Proses, Karakteristik & Komponen Fungsionalnya. *FOODREVIEW INDONESIA*, 10(8); 34–37.

Rohdiana, D. and Shabri. 2012. Aktivitas penangkapan radikal bebas DPPH (1,1-diphenyl-2-picrylhidrazyl) beberapa minuman teh dalam kemasan. *Jurnal Penelitian Teh Dan Kina*, 15(2); 89–93.

Saini, M., Sharma, M. and Kaur, L. 2016. Development of Low Sugar Green Tea Kiwi Jam from Green Tea Extract and Kiwi Pulp. *WJPLS*, 2(3); 281–287.

Scharbert, S. and Hofmann, T. 2005. Molecular definition of black tea taste by means of quantitative studies, taste reconstitution, and omission experiments. *J. Agric. Food Chem.*, 53, 5377–5384.

Senanayake, S. P. J. N. 2013. Green tea extract: Chemistry, antioxidant properties and food applications – A review. *Journal of Functional Foods*.

Shebis, Y., Iluz, D., Kinel-Tahan, Y., Dubinsky, Z. and Yehoshua, Y. 2013. Natural Antioxidants: Function and Sources. *Food and Nutrition Sciences*, 4, 643–649.

Shekhar, S. and Prasad, M. 2015. Comparative analysis of antioxidant properties of jasmine species by hydrogen peroxide assay. *European Journal of Biotechnology and Bioscience*, 3(2); 26–29.

Snyder, M., Clum, L. and Zulaica, A. V. 2015. “The Matcha Miracle”. Berkeley: Ulysses Press.

Spence, C. 2015. The psychological impact of food colour. *Flavour*, 4(21); 1–16.

Sultana, T., Stecher, G., Mayer, R., Trojer, L., Qureshi, M.N. and Abel, G. 2008. Quality assessment and quantitative analysis of flavonoids from tea samples of different origins by HPLC-DAD-ESI-MS. *Jornal Agricultural Food Chemistry*, 56(10); 3444-53.

Stawski, B. and Seideman, R. 2008. Confectinery products having tea. U.S. patent. 20080026099 A1.

- Tejero, J., Gayoso, S., Caro, I., Cordoba-Diaz, D., Mateo, J., Basterrechea, J. E., Jiménez, P. 2014. Comparative Analysis of the Antioxidant and Free-Radical Scavenging Activities of Different Water-Soluble Extracts of Green, Black and Oolong Tea Samples. *Food and Nutrition Sciences*, 5, 2157–2166.
- Theppakorn, T. 2016. Stability and chemical changes of phenolic compounds during Oolong tea processing. *International Food Research Journal*, 23(2); 564–574.
- Tokunaga, M. 2016. “Green Tea”. New York: Kodansha America.
- Topuz, A., Dincer, C., Torun, M., Tontul, I., Sahin-Nadeem, H., Haznedar, A., and Özdemir, F. 2014. Physicochemical properties of Turkish green tea powder: effects of shooting period, shading, and clone. *Turkish Journal of Agriculture and Forestry*, 38, 233–241.
- Vaclavik, V. A., & Christian, E. W. 2014. *Essentials of Food Science*. 4th ed. New York: Springer.
- Verma, I and Gupta, R. K. 2005. Estimation of phytochemical, nutritional, antioxidant and antibacterial activity of dried fruit of sacred figs (*Ficus religiosa*) and formulation of value added product (Hard Candy). *J Pharmacogn Phytochem*, 4(3); 257-267.
- Wahlqvist, M. L. 2013. Antioxidant relevance to human health. *Asia Pac J Clin Nutr*, 22(2); 171–176.
- Wijaya, C. H., Halimah, Kindly and Taqi, F.M. 2002. Penemu komposisi permen cajuput untuk pelega tenggorokan. Institut Pertanian Bogor. Paten Indonesia ID 0000385S.
- Wijaya, C. H., Idham, F.N. and Boy, M.B. 2011. Daya hambat minyak kayu putih dan flavour “cajuput candy” terhadap akumulasi biofilm *Streptococcus mutans* dan *Streptococcus sobrinus* secara in vitro. International Food Conference: Life Improvement through Food Technology, 1-87.
- Xu, J. Z., Yeung, S. Y. V., Chang, Q., Huang, Y. and Chen, Z.-Y. 2004. Comparison of antioxidant activity and bioavailability of tea epicatechins with their epimers. *British Journal of Nutrition*, 91, 873–881.
- Yagi, C., Ikeda, N. and Sato, D. 2010. Characteristics of Eight Japanese Tea Cultivars. *Fruits and Nuts*. Manoa.