

REFERENCES

- [1] P. F. W. M. Rosier *et al.*, “International Continence Society Good Urodynamic Practices and Terms 2016: Urodynamics, uroflowmetry, cystometry, and pressure-flow study,” *Neurourol Urodyn*, vol. 36, no. 5, pp. 1243–1260, Jun. 2017, doi: 10.1002/nau.23124.
- [2] W. Schäfer *et al.*, “Good urodynamic practices: Uroflowmetry, filling cystometry, and pressure-flow studies**,” *Neurourol Urodyn*, vol. 21, no. 3, pp. 261–274, 2002, doi: 10.1002/nau.10066.
- [3] D. K. Gupta, S. N. Sankhwar, and A. Goel, “Uroflowmetry Nomograms for Healthy Children 5 to 15 Years Old,” *Journal of Urology*, vol. 190, no. 3, pp. 1008–1014, Sep. 2013, doi: 10.1016/j.juro.2013.03.073.
- [4] A. E. MacNeily, “The Role of Uroflowmetry in the Diagnosis of Lower Urinary Tract Disorders in Children,” *Journal of Urology*, vol. 187, no. 6, pp. 1960–1961, Jun. 2012, doi: 10.1016/j.juro.2012.03.033.
- [5] M. L. Capitanucci, A. Marciano, A. Zaccara, E. la Sala, G. Mosiello, and M. de Gennaro, “Long-Term Bladder Function Followup in Boys with Posterior Urethral Valves: Comparison of Noninvasive vs Invasive Urodynamic Studies,” *Journal of Urology*, vol. 188, no. 3, pp. 953–957, Sep. 2012, doi: 10.1016/j.juro.2012.04.121.
- [6] S.-J. Chang, S. S.-D. Yang, and I.-N. Chiang, “Large voided volume suggestive of abnormal uroflow pattern and elevated post-void residual urine,” *Neurourol Urodyn*, vol. 30, no. 1, pp. 58–61, Jan. 2011, doi: 10.1002/nau.20901.
- [7] A. Gammie and M. J. Drake, “The fundamentals of uroflowmetry practice, based on International Continence Society good urodynamic practices recommendations,” *Neurourol Urodyn*, vol. 37, no. S6, pp. S44–S49, Aug. 2018, doi: 10.1002/nau.23777.
- [8] S.-J. Chang, S. S.-D. Yang, and I.-N. Chiang, “Large voided volume suggestive of abnormal uroflow pattern and elevated post-void residual urine,” *Neurourol Urodyn*, vol. 30, no. 1, pp. 58–61, Jan. 2011, doi: 10.1002/nau.20901.
- [9] L. E. Selman, C. A. Ochieng, A. L. Lewis, M. J. Drake, and J. Horwood, “Recommendations for conducting invasive urodynamics for men with lower urinary tract symptoms: Qualitative interview findings from a large randomized controlled trial (UPSTREAM),” *Neurourol Urodyn*, vol. 38, no. 1, pp. 320–329, Jan. 2019, doi: 10.1002/nau.23855.

- [10] B. von GARRELTS, “Micturition in the normal male.,” *Acta Chir Scand*, vol. 114, no. 3, pp. 197–210, Mar. 1958.
- [11] D. Hitt, K. Zvarova, and P. Zvara, “Urinary Flow Measurements Via Acoustic Signatures with Application to Telemedicine,” in *39th AIAA Fluid Dynamics Conference*, Jun. 2009. doi: 10.2514/6.2009-4300.
- [12] K. Zvarova, M. Ursiny, T. Giebink, K. Liang, J. G. Blaivas, and P. Zvara, “Recording urinary flow and lower urinary tract symptoms using sonouroflowmetry,” *Canadian Journal of Urology*, vol. 18, no. 3, pp. 5689–5694, Jun. 2011.
- [13] K. A. Backman, “Urinary flow during micturition in normal women.,” *Acta Chir Scand*, vol. 130, no. 4, pp. 357–70, Oct. 1965.
- [14] J. Gierup, “Micturition Studies in Infants and Children: *Normal Urinary Flow*,” *Scand J Urol Nephrol*, vol. 4, no. 3, pp. 191–207, Jan. 1970, doi: 10.3109/00365597009137597.
- [15] P. F. W. M. Rosier *et al.*, “International Continence Society Good Urodynamic Practices and Terms 2016: Urodynamics, uroflowmetry, cystometry, and pressure-flow study,” *Neurourol Urodyn*, vol. 36, no. 5, pp. 1243–1260, Jun. 2017, doi: 10.1002/nau.23124.
- [16] B. T. Haylen, D. Ashby, J. R. Sutherst, M. I. Frazer, and C. R. West, “Maximum and Average Urine Flow Rates in Normal Male and Female Populations-the Liverpool Nomograms,” *Br J Urol*, vol. 64, no. 1, pp. 30–38, Jul. 1989, doi: 10.1111/j.1464-410X.1989.tb05518.x.
- [17] S. Malde *et al.*, “Systematic Review of the Performance of Noninvasive Tests in Diagnosing Bladder Outlet Obstruction in Men with Lower Urinary Tract Symptoms,” *Eur Urol*, vol. 71, no. 3, pp. 391–402, Mar. 2017, doi: 10.1016/j.eururo.2016.09.026.
- [18] E. Bright, R. Pearcy, and P. Abrams, “Ultrasound estimated bladder weight in men attending the uroflowmetry clinic,” *Neurourol Urodyn*, vol. 30, no. 4, pp. 583–586, Apr. 2011, doi: 10.1002/nau.21049.
- [19] C. Gratzke *et al.*, “EAU guidelines on the assessment of non-neurogenic male lower urinary tract symptoms including benign prostatic obstruction,” *Eur Urol*, vol. 67, no. 6, pp. 1099–1109, Jun. 2015, doi: 10.1016/J.EURURO.2014.12.038.
- [20] K. Koiso, R. Nemoto, and M. Ohtani, “Urophonographic studies of benign prostatic hypertrophy,” *Journal of Urology*, vol. 145, no. 5, 1991, doi: 10.1016/S0022-5347(17)38537-3.
- [21] J. Krhut *et al.*, “Comparison between uroflowmetry and sonouroflowmetry in recording of urinary flow in healthy men,” *International Journal of Urology*, vol. 22, no. 8, pp. 761–765, Aug. 2015, doi: 10.1111/iju.12796.

- [22] Y. J. Lee, M. M. Kim, S. H. Song, and S. Lee, “A Novel Mobile Acoustic Uroflowmetry: Comparison With Contemporary Uroflowmetry,” *Int Neurourol J*, vol. 25, no. 2, pp. 150–156, Jun. 2021, doi: 10.5213/inj.2040250.125.
- [23] E. el Helou, J. Naba, K. Youssef, G. Mjaess, G. Sleilaty, and S. Helou, “Mobile sonouroflowmetry using voiding sound and volume,” *Sci Rep*, vol. 11, no. 1, p. 11250, Dec. 2021, doi: 10.1038/s41598-021-90659-9.
- [24] S. H. Song *et al.*, “Comparison of the conventional uroflowmetry and a novel mobile acoustic uroflowmetry in children: A pilot study,” Texas, 2020.
- [25] E. J. Aslim *et al.*, “Pilot study for the comparison of machine-learning augmented audio-uroflowmetry with standard uroflowmetry in healthy men,” *BMJ Innov*, vol. 6, no. 4, pp. 199–203, Oct. 2020, doi: 10.1136/bmjjinnov-2019-000382.
- [26] J. Krhut *et al.*, “Comparison between uroflowmetry and sonouroflowmetry in recording of urinary flow in healthy men,” *International Journal of Urology*, vol. 22, no. 8, pp. 761–765, Aug. 2015, doi: 10.1111/iju.12796.
- [27] P. Hurtik, M. Burda, J. Krhut, P. Zvara, and L. Lunacek, “Automatic diagnosis of voiding dysfunction from sound signal,” in *Proceedings - 2015 IEEE Symposium Series on Computational Intelligence, SSCI 2015*, 2015. doi: 10.1109/SSCI.2015.190.
- [28] D. A. Clifton, K. E. Niehaus, P. Charlton, and G. W. Colopy, “Health Informatics via Machine Learning for the Clinical Management of Patients,” *Yearb Med Inform*, vol. 10, no. 1, pp. 38–43, Aug. 2015, doi: 10.15265/IY-2015-014.
- [29] A. Rajkomar, J. Dean, and I. Kohane, “Machine Learning in Medicine,” *New England Journal of Medicine*, vol. 380, no. 14, pp. 1347–1358, 2019, doi: 10.1056/nejmra1814259.
- [30] A. Rajkomar, J. Dean, and I. Kohane, “Machine Learning in Medicine,” *New England Journal of Medicine*, vol. 380, no. 14, pp. 1347–1358, Apr. 2019, doi: 10.1056/NEJMra1814259.
- [31] S. P. R. Baas, “A machine learning approach to the automatic classification of female uroflowmetry measurements. PhD Thesis (University of Twente),” 2016.
- [32] W. Schfer *et al.*, “Good Urodynamic Practices: Uroflowmetry, filling cystometry, and pressure-flow studies,” *Neurourol Urodyn*, vol. 21, no. 3, pp. 261–274, 2002, doi: 10.1002/nau.10066.
- [33] S. P. R. Baas, “A machine learning approach to the automatic classification of female uroflowmetry measurements.” Jun. 28, 2016. [Online]. Available: <http://essay.utwente.nl/70570/>

- [34] Y. Li, W. Li, J. Xiong, J. Xia, and Y. Xie, “Comparison of Supervised and Unsupervised Deep Learning Methods for Medical Image Synthesis between Computed Tomography and Magnetic Resonance Images,” *Biomed Res Int*, vol. 2020, p. 5193707, 2020, doi: 10.1155/2020/5193707.
- [35] M. Alloghani, D. Al-Jumeily, J. Mustafina, A. Hussain, and A. J. Aljaaf, “A systematic review on supervised and unsupervised machine learning algorithms for data science,” *Supervised and unsupervised learning for data science*, pp. 3–21, 2020.
- [36] A. Albalate and W. Minker, *Semi-supervised and unsupervised machine learning: novel strategies*. John Wiley & Sons, 2013.
- [37] Q. K. Al-Shayea, “Artificial neural networks in medical diagnosis,” *International Journal of Computer Science Issues*, vol. 8, no. 2, pp. 150–154, 2011.
- [38] Z. R. Yang and Z. Yang, “Artificial Neural Networks,” *Comprehensive Biomedical Physics*, vol. 6, pp. 1–17, 2014, doi: 10.1016/B978-0-444-53632-7.01101-1.
- [39] P. Rosier, “Contemporary diagnosis of lower urinary tract dysfunction.,” *F1000Res*, vol. 8, p. 644, May 2019, doi: 10.12688/f1000research.16120.1.
- [40] M. J. Drake, S. K. Doumouchtsis, H. Hashim, and A. Gammie, “Fundamentals of urodynamic practice, based on International Continence Society good urodynamic practices recommendations,” *Neurourol Urodyn*, vol. 37, no. S6, pp. S50–S60, Aug. 2018, doi: 10.1002/nau.23773.
- [41] M. B. Siroky, C. A. Olsson, and R. J. Krane, “The Flow Rate Nomogram: II. Clinical Correlation,” *Journal of Urology*, vol. 123, no. 2, pp. 208–210, Feb. 1980, doi: 10.1016/S0022-5347(17)55859-0.
- [42] C. Gratzke *et al.*, “EAU Guidelines on the Assessment of Non-neurogenic Male Lower Urinary Tract Symptoms including Benign Prostatic Obstruction,” *Eur Urol*, vol. 67, no. 6, pp. 1099–1109, Jun. 2015, doi: 10.1016/j.eururo.2014.12.038.
- [43] G. S. Sonke, C. Robertson, A. L. M. Verbeek, W. P. J. Witjes, J. J. M. C. H. de la Rosette, and L. A. Kiemeney, “A method for estimating within-patient variability in maximal urinary flow rate adjusted for voided volume,” *Urology*, vol. 59, no. 3, pp. 368–372, Mar. 2002, doi: 10.1016/S0090-4295(01)01547-3.
- [44] G. S. Sonke, C. Robertson, A. L. M. Verbeek, W. P. J. Witjes, J. J. M. C. H. de la Rosette, and L. A. Kiemeney, “A method for estimating within-patient variability in maximal urinary flow rate adjusted for voided volume,” *Urology*, vol. 59, no. 3, pp. 368–372, Mar. 2002, doi: 10.1016/S0090-4295(01)01547-3.

- [45] M. R. Feneley, W. D. Dunsmuir, J. Pearce, and R. S. Kirby, “Reproducibility of uroflow measurement: experience during a double-blind, placebo-controlled study of doxazosin in benign prostatic hyperplasia,” *Urology*, vol. 47, no. 5, pp. 658–663, May 1996, doi: 10.1016/S0090-4295(96)00014-3.